1 CLAIMS

2

- 3 1. A method of preparing a composition for the
- 4 treatment of dermatological disorders, wherein coal
- 5 tar is filtered to remove impurities by filter
- 6 compression.

7

- 8 2. A method as in Claim, wherein zinc pyrithione is
- 9 incorporated into the composition.

10

- 11 3. A method as in any of the previous Claims, wherein
- the coal tar is filtered by being fed through a
- 13 compressed charcoal filter.

14

- 15 4. A method as in Claim 3, wherein the coal tar is fed
- 16 through the filter at 17 to 19 psi.

17

- 18 5. A method as in Claim 3, wherein the coal tar is fed
- 19 through the filter at 18 psi.

20

- 21 6. A method as in Claims 3 to 5, wherein compressed air
- is used to force the coal tar through the filter.

23

- 24 7. A method as in Claims 3 to 6, wherein the coal tar
- is left in the filter system for 8 hours.

26

- 27 8. A method as in Claims 3 to 7, wherein the fluid that
- has been passed through the filter is boiled.

29

- 30 9. A method as in Claim 8, wherein the fluid that has
- 31 been passed through the filter is boiled for 5
- 32 minutes.

- 1 10. A method as in Claims 8 or 9, wherein the boiled,
- 2 filtered fluid is allowed to cool to room
- 3 temperature.

4

- 5 11. A method as in Claims 8, 9 or 10, wherein the top
- 6 layer of the boiled, filtered fluid is refiltered.

7

- 8 12. A method as in Claim 11, wherein refiltering is
- 9 through a nylon mesh.

10

- 11 13. A method as in any of the previous Claims, where a
- 12 surfactant is added to the formulation.

13

- 14 14. A method as in Claim 13, wherein the surfactant is
- 15 an ionic surfactant.

16

- 17 15. A method as in Claims 13 or 14, wherein the
- surfactant is sodium lauryl sulphate.

19

- 20 16. A method as in any of the previous Claims, wherein a
- 21 carrier is added to the formulation.

22

- 23 17. A method as in any of the previous Claims, wherein
- 24 multiple carriers are added to the formulation.

25

- 26 18. A method as in Claims 16 or 17, wherein the carrier
- is isopropyl myristate.

28

- 29 19. A method as in Claims 16 or 17, wherein the carrier
- 30 is ethyl alcohol.

- 1 20. A method as in any of the previous Claims, wherin
- 2 the formulation is placed in a spray or aerosol
- 3 container.

4

5 21. A composition for the treatment of dermatological disorders comprising:

7

- 6 coal tar;
- 9 zinc pyrithione;
- one or more surfactants; and
- one or more carriers

12

- 13 22. A method as in Claim 21, wherein the composition
- 14 also contains allantoin.

15

- 16 23. A composition as in Claims 21 and 22, wherein the
- 17 surfactant is an ionic surfactant.

18

- 19 24. A composition as in Claims 21 to 23, wherein the
- 20 surfactant is sodium lauryl sulphate.

21

- 22 25. A composition as in Claims 21 to 24, wherein the
- 23 carrier is isopropyl myristate.

24

- 25 26. A composition as in Claims 21 to 24, wherein the
- 26 carrier is ethyl alcohol.

27

- 28 27. A composition as in Claims 21 to 25, wherein the
- 29 composition also comprises an anti-fungal agent.

30

- 31 28. A composition as in Claim 27, wherein the anti-
- fungal agent is undecylenic acid.

1	29.	A composition for the treatment of dermatological	al
2		disorders, comprising:	
3			
4		zinc pyrithione;	
5		 alcoholic extract of coal tar; 	
6		allantoin;	
7		sodium lauryl sulphate;	
8		isopropyl myristate;	
9		ethyl alcohol; and	
10		• undecylenic acid	
11			
12	30.		
13		ingredients in the following amounts:	
14		,	
15		• ZINC pyrithione).20%
16		• alcoholic exclact of coal cal	0.25%
17		• allancoin	0.25%
18		• Soutum tautyt sutphace	0.10%
19		• isopropyl myristate 4:	9.45%
20		• ethyl alcohol 4:	9.45%
21		• undecylenic acid	0.30%
22			
23	31.	_	ne
24		composition is provided in a spray form.	
25			ho
26	32.	-	пе
27		composition is provided in an aerosol form.	